

Classifications of some classes of Zinbiel algebras.

J.Q. Adashev¹

*Department of Mathematics, National University of Uzbekistan, Tashkent,
Uzbekistan*

e-mails: adashevjq@mail.ru

B.A. Omirov²

Institute of Mathematics, Uzbek Academy of Sciences, Tashkent, Uzbekistan

e-mail: omirovb@mail.ru

A.Kh. Khudoyberdiyev

*Department of Mathematics, National University of Uzbekistan, Tashkent,
Uzbekistan*

e-mail: abror_alg_85@mail.ru

Abstract

In this work nul-filiform and filiform Zinbiel algebras are described up to isomorphism. Moreover, the classification of complex Zinbiel algebras is extended from dimensions ≤ 3 up to the dimension 4.

Key words: Zinbiel algebra, nilpotency, nul-filiform algebra, filiform algebra, Leibniz algebra.

MR(2000) Subject Classification: 17A32.

1 Introduction

One of the important object of the modern theory of non-associative algebras is Lie algebras. Active investigations in the theory of Lie algebras lead to the appearing of some generalizations of these algebras such as Mal'cev algebras, Lie superalgebras, binary Lie algebras, Leibniz algebras and others.

In the present work, we consider algebras which are dual to Leibniz algebras. Recall that Leibniz algebras were introduced in [6] in the 90-th years of the last century. They are defined by the following identity

$$[x, [y, z]] = [[x, y], z] - [[x, z], y].$$

¹ This work is supported by ICTP

² and DFG 436 USB 113/10/0-1 project (Germany).

J.-L. Loday in [5] studied categorical properties of Leibniz algebras and considered in this connection a new object — Zinbiel algebra (read Leibniz in reverse order). Since the category of Zinbiel algebras is Koszul dual to the category of Leibniz algebras, sometimes they are also called dual Leibniz algebras [7].

In works [3]–[4] some interesting properties of Zinbiel algebras were obtained. In particular, the nilpotency of an arbitrary complex finite dimensional Zinbiel algebra was proved in [4].

For the examples of Zinbiel algebras we refer to works [4], [5] and [7].

Since description of all finite dimensional complex Zinbiel algebras (which are nilpotent) is a boundless problem, it is natural to add certain restrictions for their investigation. One of such restriction is restriction on the nilindex. Note works [2] and [8], where structural description of nilpotent Lie algebras and Leibniz algebras is given.

In study of any class of algebras, it is important to describe up to isomorphism even algebras of lower dimensions because such description gives examples for to establish or reject certain conjectures. In this way in [4], the classification of complex Zinbiel algebras of dimensions ≤ 3 is given. Applying some general results obtained for finite dimensional Zinbiel algebras we extended classification of complex Zinbiel algebras up to dimension 4.

2 Classification of complex nul-filiform Zinbiel algebras

Definition 2.1. *An algebra A over a field F is called Zinbiel algebra if for any $x, y, z \in A$ the following identity*

$$(x \circ y) \circ z = x \circ (y \circ z) + x \circ (z \circ y) \quad (1)$$

holds.

For a given Zinbiel algebra A we define the following sequence:

$$A^1 = A, A^{k+1} = A \circ A^k, k \geq 1.$$

Definition 2.2. *An Zinbiel algebra A is called nilpotent if there exists $s \in \mathbb{N}$ such that $A^s = 0$. The minimal number s satisfying this property is called index of nilpotency or nilindex of the algebra A .*

It is not difficult to see that the index of nilpotency of an arbitrary n -dimensional nilpotent algebra does not exceed the number $n + 1$.

Definition 2.3. *An n -dimensional Zinbiel algebra A is called nul-filiform if $\dim A^i = (n + 1) - i$, $1 \leq i \leq n + 1$.*

It is evident, the last definition is equivalent to the fact that algebra A has maximal index of nilpotency.

Theorem 2.1. *An arbitrary n -dimensional nul-filiform Zinbiel algebra is isomorphic to the algebra:*

$$e_i \circ e_j = C_{i+j-1}^j e_{i+j}, \quad \text{for } 2 \leq i + j \leq n \quad (2)$$

where omitted products are equal to zero and $\{e_1, e_2, \dots, e_n\}$ is a basis of the algebra.

Proof. Let A be an n -dimensional nul-filiform Zinbiel algebra and let $\{x_1, x_2, \dots, x_n\}$ is a basis of the algebra A such that $x_1 \in A^1 \setminus A^2$, $x_2 \in A^2 \setminus A^3$, $\dots$, $x_n \in A^n$. Since $x_2 \in A^2 \setminus A^3$, we have for some elements $b_{2,p}, c_{2,p}$ of algebra A

$$x_2 = \sum b_{2,p} \circ c_{2,p} = \sum \alpha_{i,j} x_i \circ x_j = \alpha_{1,1} x_1 \circ x_1 + (*)$$

where $(*) \in A^3$, i.e. $x_2 = \alpha_{1,1} x_1 \circ x_1 + (*)$. Note that $\alpha_{1,1} x_1 \circ x_1 \neq 0$. Indeed, in the opposite case $x_2 \in A^3$.

Similar, for $x_3 \in A^3 \setminus A^4$ we have

$$x_3 = \sum a_{3,p} \circ (b_{3,p} \circ c_{3,p}) = \sum \alpha_{i,j,k} x_i \circ (x_j \circ x_k) = \alpha_{1,1,1} x_1 \circ (x_1 \circ x_1) + (**)$$

where $(**) \in A^4$ and $\alpha_{1,1,1} x_1 \circ (x_1 \circ x_1) \neq 0$ (otherwise $x_3 \in A^4$), i.e. $x_3 = \alpha_{1,1,1} x_1 \circ (x_1 \circ x_1) + (**)$. Continuing this procedure we obtain that elements $e_1 := x_1$, $e_2 := x_1 \circ x_1$, $e_3 := x_1 \circ (x_1 \circ x_1)$, $\dots$, $e_n := (x_1 \circ \dots \circ (x_1 \circ (x_1 \circ x_1)))$ are different from zero. It is not difficult to check the linear independence of these elements. Hence we can choose as a basis of algebra A the elements $\{e_1, e_2, \dots, e_n\}$. We have by construction

$$e_1 \circ e_i = e_{i+1} \quad \text{for } 1 \leq i \leq n - 1. \quad (3)$$

We shall prove the equality (2) by induction on j for any i .

Using identities (1), (3) we can prove by induction the equality:

$$e_i \circ e_1 = i e_{i+1} \quad \text{for } 1 \leq i \leq n - 1,$$

i.e. equality (2) is true for $j = 1$ and any i .

Suppose that the equality is true for all $j \leq k - 1$ and any i .

Let us prove the equality (2) for $j = k$ and any i . Using the inductive hypothesis and the following chain of equalities:

$$\begin{aligned} e_i \circ e_k &= e_i \circ (e_1 \circ e_{k-1}) = (e_i \circ e_1) \circ e_{k-1} - e_i \circ (e_{k-1} \circ e_1) = \\ &= ie_{i+1} \circ e_{k-1} - (k-1)e_i \circ e_k = iC_{i+k-1}^{k-1}e_{i+k} - (k-1)e_i \circ e_k \end{aligned}$$

we obtain $ke_i \circ e_k = iC_{i+k-1}^{k-1}e_{i+k}$, i.e.

$$e_i \circ e_k = \frac{i}{k}C_{i+k-1}^{k-1}e_{i+k} = \frac{i}{k} \frac{(i+k-1)!}{(k-1)!i!}e_{i+k} = \frac{(i+k-1)!}{k!(i-1)!}e_{i+k} = C_{i+k-1}^k e_{i+k}.$$

□

We denote the algebra from theorem 2.1 as NF_n .

It is not difficult to see that n -dimensional Zinbiel algebra is one generated if and only if it is isomorphic to the algebra NF_n .

3 Classification of complex filiform Zinbiel algebras

Definition 3.1. An n -dimensional Zinbiel algebra A is said to be filiform if $\dim A^i = n - i$, $2 \leq i \leq n$.

Consider the natural gradation of an filiform Zinbiel algebra taking $A_i = A^i/A^{i+1}$, $1 \leq i \leq n - 1$. It is obvious that $\dim A_1 = 2$ and $\dim A_i = 1$, $2 \leq i \leq n - 1$, and

$$A \cong A_1 \oplus A_2 \oplus \dots \oplus A_{n-1} \quad \text{where } A_i \circ A_j \subseteq A_{i+j}.$$

In the following theorem, the classification of complex naturally graded filiform Zinbiel algebras is represented.

Theorem 3.1. An arbitrary n -dimensional ($n \geq 5$) naturally graded complex filiform Zinbiel algebra is isomorphic to the following algebra:

$$F_n : e_i \circ e_j = C_{i+j-1}^j e_{i+j}, \quad 2 \leq i+j \leq n-1 \quad (4)$$

where omitted products are equal to zero and $\{e_1, e_2, \dots, e_n\}$ is a basis of the algebra.

Proof. Let A be an Zinbiel algebra satisfying conditions of the theorem. Similar to the work [8] we take a basis $\{e_1, e_2, \dots, e_n\}$ of the algebra A such that $A_1 = \langle e_1, e_n \rangle$, $A_2 = \langle e_2 \rangle$, $A_3 = \langle e_3 \rangle$, $\dots$, $A_{n-1} = \langle e_{n-1} \rangle$, and $e_1 \circ e_i = e_{i+1}$ at $2 \leq i \leq n-2$.

Introduce the following notations

$$e_1 \circ e_1 = \alpha_1 e_2, \quad e_1 \circ e_n = \alpha_2 e_2, \quad e_n \circ e_1 = \alpha_3 e_2, \quad e_n \circ e_n = \alpha_4 e_2.$$

Now consider the possible cases.

Case 1. Let $(\alpha_1, \alpha_4) \neq (0, 0)$. Then without any loss of generality we can suppose $\alpha_1 \neq 0$. Change the basis as follows $e'_2 = \alpha_1 e_2$, $e'_3 = \alpha_1 e_3$, $\dots$, $e'_{n-1} = \alpha_1 e_{n-1}$ we can suppose $\alpha_1 = 1$.

In this case, the space spanned on vectors $\{e_1, e_2, \dots, e_{n-1}\}$ forms a nilform Zinbiel algebra of the dimension $n-1$. From the proof of theorem 2.1 we can conclude $e_i \circ e_j = C_{i+j-1}^j e_{i+j}$, $2 \leq i+j \leq n-1$.

Consider identity (1) in the following multiplications:

$$(e_1 \circ e_n) \circ e_1 = e_1 \circ (e_n \circ e_1) + e_1 \circ (e_1 \circ e_n) \Rightarrow 2\alpha_2 e_3 = \alpha_3 e_3 + \alpha_2 e_3,$$

i.e. $\alpha_2 = \alpha_3$;

$$(e_1 \circ e_1) \circ e_n = e_1 \circ (e_1 \circ e_n) + e_1 \circ (e_n \circ e_1) \Rightarrow e_2 \circ e_n = \alpha_3 e_3 + \alpha_2 e_3,$$

hence, $e_2 \circ e_n = 2\alpha_2 e_3$;

$$(e_1 \circ e_n) \circ e_n = 2e_1 \circ (e_n \circ e_n) \Rightarrow 2\alpha_2^2 e_3 = 2\alpha_4 e_3,$$

consequently, $\alpha_2^2 = \alpha_4$;

$$(e_n \circ e_1) \circ e_1 = 2e_n \circ (e_1 \circ e_1) \Rightarrow 2\alpha_2 e_3 = 2e_n \circ e_2,$$

i.e. we have $e_n \circ e_2 = \alpha_2 e_3$.

Take the change of the basic elements by the following way:

$$e'_n = -\alpha_2 e_1 + e_n, \quad e'_i = e_i \text{ for } 1 \leq i \leq n-1.$$

It is not difficult to see that $e'_1 \circ e'_n = e'_n \circ e'_1 = e'_n \circ e'_n = e'_2 \circ e'_n = e'_n \circ e'_2 = 0$, moreover other products are not changed, i.e. we can suppose $\alpha_2 = \alpha_3 = \alpha_4 = 0$.

Using identity (1) and the method of mathematical induction one can easy to prove

$$e_n \circ e_i = 0 \text{ for } 1 \leq i \leq n-1. \quad (5)$$

The equality

$$e_i \circ e_n = 0, \text{ for } 1 \leq i \leq n-1$$

can be proved by induction using (1) and (5).

Case 2. Let $(\alpha_1, \alpha_4) = (0, 0)$. Then $(\alpha_2, \alpha_3) \neq (0, 0)$. In the case of $\alpha_2 \neq -\alpha_3$ we obtain case 1 taking $e'_1 = e_1 + e_n$. Now consider the case $\alpha_2 = -\alpha_3 \neq 0$. Taking the following change of basis

$$e'_1 = e_1, \quad e'_n = e_n, \quad e'_i = \alpha_2 e_i \text{ for } 2 \leq i \leq n-1$$

we can suppose $\alpha_2 = 1$.

Consider the following products

$$(e_1 \circ e_n) \circ e_1 = e_1 \circ (e_n \circ e_1) + e_1 \circ (e_1 \circ e_n) \Rightarrow e_2 \circ e_1 = 0;$$

$$0 = (e_1 \circ e_1) \circ e_2 = e_1 \circ (e_1 \circ e_2) + e_1 \circ (e_2 \circ e_1) = e_1 \circ e_3 = e_4.$$

Thus, we obtain the contradiction with existence of an algebra in this case. $\square$

The following proposition allows to extract a "convenient" basis in an arbitrary complex filiform Zinbiel algebra. Such basis is often called in literature adapted [8].

Proposition 3.1. *There exists basis $\{e_1, e_2, \dots, e_n\}$ in an arbitrary n -dimensional ($n \geq 5$) complex filiform Zinbiel algebra such that the multiplication of the algebra has the following form:*

$$\begin{aligned} e_i \circ e_j &= C_{i+j-1}^j e_{i+j}, \quad 2 \leq i+j \leq n-1 \\ e_n \circ e_1 &= \alpha e_{n-1}, \quad e_n \circ e_n = \beta e_{n-1} \end{aligned} \tag{6}$$

where $\alpha, \beta \in \mathbb{C}$.

Proof. By theorem 3.1 we have that any n -dimensional complex filiform Zinbiel algebra is isomorphic to an algebra of the form:

$$F_n + \beta$$

where

$$\begin{aligned} \beta(e_1, e_i) &= 0, \quad \text{for } 1 \leq i \leq n-1, \\ \beta(e_n, e_n) &\in \text{lin}\{e_3, e_4, \dots, e_{n-1}\}, \end{aligned}$$

$$\beta(e_i, e_n), \beta(e_n, e_i) \in \text{lin}\{e_{i+2}, e_{i+3}, \dots, e_{n-1}\} \text{ for } 1 \leq i \leq n-1,$$

$$\beta(e_i, e_1) \in \text{lin}\{e_{i+2}, e_{i+3}, \dots, e_{n-1}\} \text{ for } 2 \leq i \leq n-1,$$

$$\beta(e_i, e_j) \in \text{lin}\{e_{i+j+1}, e_{i+j+2}, \dots, e_{n-1}\} \text{ for } 2 \leq i \leq n-1.$$

Similarly as in proof of the theorem 3.1 it is not difficult to establish that the multiplications:

$$e_i \circ e_1, \quad e_i \circ e_j, \quad 1 \leq i, j \leq n-1$$

can be obtained from $e_1 \circ e_i = e_{i+1}$, $1 \leq i \leq n-2$ and identity (1).

By the similar procedure, we obtain

$$\beta(e_i, e_1) = 0, \text{ for } 1 \leq i \leq n-1,$$

$$\beta(e_i, e_j) = 0, \text{ for } 2 \leq i, j \leq n-1.$$

Let a Zinbiel algebra A be isomorphic to the algebra $F_n + \beta$. Then we have

$$e_1 \circ e_i = e_{i+1}, \text{ for } 1 \leq i \leq n-2.$$

Put $e_1 \circ e_n = \alpha_3 e_3 + \alpha_4 e_4 + \dots + \alpha_{n-1} e_{n-1}$, $e_n \circ e_1 = \beta_3 e_3 + \beta_4 e_4 + \dots + \beta_{n-1} e_{n-1}$.

Taking the change $e'_n = e_n - \alpha_3 e_2 - \alpha_4 e_3 - \dots - \alpha_{n-1} e_{n-2}$, we can suppose $\alpha_3 = \alpha_4 = \dots = \alpha_{n-1} = 0$, i.e. $e_1 \circ e_n = 0$.

Identity (1) implies

$$(e_1 \circ e_n) \circ e_1 = e_1 \circ (e_n \circ e_1) + e_1 \circ (e_1 \circ e_n) \Rightarrow e_1 \circ (\beta_3 e_3 + \beta_4 e_4 + \dots + \beta_{n-1} e_{n-1}) = 0$$

therefore $\beta_3 e_4 + \beta_4 e_5 + \dots + \beta_{n-2} e_{n-1} = 0$. Hence, $e_n \circ e_1 = \beta_{n-1} e_{n-1}$.

Consider the product

$$(e_n \circ e_n) \circ e_1 = e_n \circ (e_n \circ e_1) + e_n \circ (e_1 \circ e_n) = 0$$

from which we obtain $e_n \circ e_n = \gamma e_{n-1}$ for some γ .

Similarly to the proof of theorem 3.1 we can obtain that $e_n \circ e_i = 0$ for $2 \leq i \leq n-1$ and $e_i \circ e_n = 0$ for $1 \leq i \leq n-1$.

Thus, we obtain that Zinbiel algebra A is split, i.e. $A = NF_{n-1} \oplus C$. Applying theorem 2.1 complete the proof of the proposition. $\square$

The classification of complex filiform Zinbiel algebras is given in the following

Theorem 3.2. *An arbitrary n -dimensional ($n \geq 5$) complex filiform Zinbiel algebras is isomorphic to one of the following pairwise non isomorphic algebras:*

$$\begin{aligned} F_n^1 : e_i \circ e_j &= C_{i+j-1}^j e_{i+j}, \quad 2 \leq i+j \leq n-1; \\ F_n^2 : e_i \circ e_j &= C_{i+j-1}^j e_{i+j}, \quad 2 \leq i+j \leq n-1, \quad e_n \circ e_1 = e_{n-1}; \\ F_n^3 : e_i \circ e_j &= C_{i+j-1}^j e_{i+j}, \quad 2 \leq i+j \leq n-1, \quad e_n \circ e_n = e_{n-1}. \end{aligned}$$

Proof. By proposition 3.1 we have the multiplication (multiplication (6)) in an n -dimensional complex filiform Zinbiel algebra, namely,

$$\begin{aligned} e_i \circ e_j &= C_{i+j-1}^j e_{i+j}, \quad 2 \leq i+j \leq n-1 \\ e_n \circ e_1 &= \alpha e_{n-1}, \quad e_n \circ e_n = \beta e_{n-1}. \end{aligned}$$

Consider the general change of generator basic elements in the form:

$$\begin{aligned} e'_1 &= a_1 e_1 + a_2 e_2 + \dots + a_n e_n, \\ e'_n &= b_1 e_1 + b_2 e_2 + \dots + b_n e_n, \end{aligned}$$

where $a_1 \neq 0$ and $a_1 b_n - a_n b_1 \neq 0$.

Then expressing the rest basic elements of the new basis by means of basis elements of the old basis and comparing the obtained relations, we obtain the following restrictions:

$$\begin{aligned} a_1 b_1 &= 0, \\ a_1 b_2 + 2a_2 b_1 &= 0, \\ a_1 b_3 + C_3^2 a_2 b_2 + 3a_3 b_1 &= 0, \\ &\dots\dots\dots, \\ a_1 b_{n-3} + C_{n-3}^{n-4} a_2 b_{n-4} + C_{n-3}^{n-5} a_3 b_{n-5} + \dots + (n-3)a_{n-3} b_1 &= 0, \\ a_1 b_{n-2} + C_{n-2}^{n-3} a_2 b_{n-3} + C_{n-2}^{n-4} a_3 b_{n-4} + \dots + (n-2)a_{n-2} b_1 + \alpha a_n b_1 + \beta a_n b_n &= 0. \end{aligned}$$

We have recurrently from these restrictions $b_1 = b_2 = \dots = b_{n-3} = 0$. Hence, $b_{n-2} = -\frac{a_n b_n}{a_1} \beta$.

Consider the product

$$e'_n \circ e'_1 = \left(-\frac{a_n b_n}{a_1} \beta e_{n-2} + b_{n-1} e_{n-1} + b_n e_n\right) \circ (a_1 e_1 + a_2 e_2 + \dots + a_n e_n) =$$

$$= -(n-2)\beta a_n b_n e_{n-1} + \alpha a_1 b_n e_{n-1} + \beta a_n b_n e_{n-1} = (\alpha a_1 b_n - (n-3)a_n b_n) e_{n-1}.$$

On the other hand,

$$e'_n \circ e'_1 = \alpha' a_1^{n-1} e_{n-1}.$$

Comparing coefficients at e_{n-1} we obtain

$$\alpha a_1 b_n - (n-3)\beta a_n b_n = \alpha' a_1^{n-1}.$$

Consider the product

$$e'_n \circ e'_n = \left(-\frac{a_n b_n}{a_1} \beta e_{n-2} + b_{n-1} e_{n-1} + b_n e_n\right) \circ \left(-\frac{a_n b_n}{a_1} \beta e_{n-2} + b_{n-1} e_{n-1} + b_n e_n\right) = b_n^2 \beta e_{n-1}.$$

On the other hand,

$$e'_n \circ e'_n = \beta' a_1^{n-1} e_{n-1}.$$

Comparing coefficients we obtain

$$b_n^2 \beta = \beta' a_1^{n-1}.$$

Now consider the following cases.

Case 1. Let $\beta = 0$. Then $\beta' = 0$ and $\alpha b_n = \alpha' a_1^{n-2}$. If $\alpha = 0$, then $\alpha' = 0$ and we have algebra F_n^1 . If $\alpha \neq 0$, then taking $b_n = \frac{a_1^{n-2}}{\alpha}$ we get $\alpha' = 1$, i.e. the algebra F_n^2 is obtained.

Case 2. Let $\beta \neq 0$. Then putting $b_n = \sqrt{a_1^{n-1}}$, $a_n = \frac{\alpha a_1}{(n-3)\beta}$ we get $\beta' = 1$, $\alpha' = 0$ and algebra F_n^3 is obtained.

Note that the obtained algebras are not pairwise isomorphic. $\square$

Comparing the description of complex filiform Leibniz algebras [2] and the result of theorem 3.2 we can note how much the class of complex filiform Zinbiel algebras is "thin". So, although Zinbiel algebras and Leibniz algebras are Koszul dual, but they quantitatively strongly are distinguished even in the class of filiform algebras .

4 Classification of four dimensional complex Zinbiel algebras.

Since an arbitrary finite dimensional complex Zinbiel algebra is nilpotent, therefore for an arbitrary four dimensional Zinbiel algebra A the condition $A^5 = 0$ holds.

As the direct sum of nilpotent Zinbiel algebras is nilpotent, then in the description it is enough to consider non split algebras.

Theorem 4.1. *An arbitrary four-dimensional complex non split Zinbiel algebra is isomorphic to the one of the following pairwise non isomorphic algebras:*

$$\begin{aligned}
A_1 : \quad & e_1 \circ e_1 = e_2, \quad e_1 \circ e_2 = e_3, \quad e_2 \circ e_1 = 2e_3, \quad e_1 \circ e_3 = e_4, \quad e_2 \circ e_2 = 3e_4, \\
& e_3 \circ e_1 = 3e_4; \\
A_2 : \quad & e_1 \circ e_1 = e_3, \quad e_1 \circ e_2 = e_4, \quad e_1 \circ e_3 = e_4, \quad e_3 \circ e_1 = 2e_4; \\
A_3 : \quad & e_1 \circ e_1 = e_3, \quad e_1 \circ e_3 = e_4, \quad e_2 \circ e_2 = e_4, \quad e_3 \circ e_1 = 2e_4; \\
A_4 : \quad & e_1 \circ e_2 = e_3, \quad e_1 \circ e_3 = e_4, \quad e_2 \circ e_1 = -e_3; \\
A_5 : \quad & e_1 \circ e_2 = e_3, \quad e_1 \circ e_3 = e_4, \quad e_2 \circ e_1 = -e_3, \quad e_2 \circ e_2 = e_4; \\
A_6 : \quad & e_1 \circ e_1 = e_4, \quad e_1 \circ e_2 = e_3, \quad e_2 \circ e_1 = -e_3, \quad e_2 \circ e_2 = -2e_3 + e_4; \\
A_7 : \quad & e_1 \circ e_2 = e_3, \quad e_2 \circ e_1 = e_4, \quad e_2 \circ e_2 = -e_3; \\
A_8(\alpha) : \quad & e_1 \circ e_1 = e_3, \quad e_1 \circ e_2 = e_4, \quad e_2 \circ e_1 = -\alpha e_3, \quad e_2 \circ e_2 = -e_4, \quad \alpha \in C; \\
A_9(\alpha) : \quad & e_1 \circ e_1 = e_4, \quad e_1 \circ e_2 = \alpha e_4, \quad e_2 \circ e_1 = -\alpha e_4, \quad e_2 \circ e_2 = e_4, \\
& e_3 \circ e_3 = e_4, \quad \alpha \in C; \\
A_{10} : \quad & e_1 \circ e_2 = e_4, \quad e_1 \circ e_3 = e_4, \quad e_2 \circ e_1 = -e_4, \quad e_2 \circ e_2 = e_4, \quad e_3 \circ e_1 = e_4; \\
A_{11} : \quad & e_1 \circ e_1 = e_4, \quad e_1 \circ e_2 = e_4, \quad e_2 \circ e_1 = -e_4, \quad e_3 \circ e_3 = e_4; \\
A_{12} : \quad & e_1 \circ e_2 = e_3, \quad e_2 \circ e_1 = e_4; \\
A_{13} : \quad & e_1 \circ e_2 = e_3, \quad e_2 \circ e_1 = -e_3, \quad e_2 \circ e_2 = e_4; \\
A_{14} : \quad & e_2 \circ e_1 = e_4, \quad e_2 \circ e_2 = e_3; \\
A_{15}(\alpha) : \quad & e_1 \circ e_2 = e_4, \quad e_2 \circ e_2 = e_3, \quad e_2 \circ e_1 = \frac{1+\alpha}{1-\alpha} e_4, \quad \alpha \in C \setminus \{1\}; \\
A_{16} : \quad & e_1 \circ e_2 = e_4, \quad e_2 \circ e_1 = -e_4, \quad e_3 \circ e_3 = e_4.
\end{aligned}$$

Proof. Note that the result of Proposition 3.1 of [1] also holds for Zinbiel algebras. Therefore we have the following possible cases for $(\dim A^2, \dim A^3, \dim A^4)$:

$$(3, 2, 1), (2, 1, 0), (2, 0, 0), (1, 0, 0), (0, 0, 0).$$

It is obvious, an Zinbiel algebra with the condition $(3, 2, 1)$ is nul-filiform. Using theorem 2.1 we obtain algebra A_1 .

Consider an algebra with the restriction $(2, 1, 0)$ (this algebra is filiform).

Let $\{e_1, e_2, e_3, e_4\}$ be a basis of algebra A satisfying the conditions $A^2 = \{e_3, e_4\}$, $A^3 = \{e_4\}$. Then we can suppose that

$$\begin{aligned}
e_1 \circ e_1 &= \alpha_1 e_3 + \alpha_2 e_4, & e_1 \circ e_2 &= \alpha_3 e_3 + \alpha_4 e_4, \\
e_2 \circ e_1 &= \alpha_5 e_3 + \alpha_6 e_4, & e_2 \circ e_2 &= \alpha_7 e_3 + \alpha_8 e_4, \\
e_1 \circ e_3 &= e_4, & e_2 \circ e_3 &= \alpha_9 e_4,
\end{aligned}$$

where $(\alpha_1, \alpha_3, \alpha_5, \alpha_7) \neq (0, 0, 0, 0)$.

Case 1. Let $(\alpha_1, \alpha_7) \neq (0, 0)$. Then by arguments analogous to arguments in the proofs of theorems 3.1 and 3.2 we obtain algebras:

$$e_1 \circ e_1 = e_3, \quad e_1 \circ e_3 = e_4, \quad e_3 \circ e_1 = 2e_4;$$

$$e_1 \circ e_1 = e_3, \quad e_1 \circ e_2 = e_4, \quad e_1 \circ e_3 = e_4, \quad e_3 \circ e_1 = 2e_4;$$

$$e_1 \circ e_1 = e_3, \quad e_1 \circ e_3 = e_4, \quad e_2 \circ e_2 = e_4, \quad e_3 \circ e_1 = 2e_4.$$

Note that the algebra defined by multiplication

$$e_1 \circ e_1 = e_3, \quad e_1 \circ e_3 = e_4, \quad e_3 \circ e_1 = 2e_4$$

is split. So, in this case we have the algebras $A_2 - A_3$.

Case 2. Let $(\alpha_1, \alpha_7) = (0, 0)$. Then $(\alpha_3, \alpha_5) \neq (0, 0)$. If $\alpha_3 \neq -\alpha_5$, then taking $e'_1 = Ae_1 + e_2$, where $A \neq -\alpha_9$, we have case 1. It remains to consider the case $\alpha_3 = -\alpha_5$. Denote $e'_3 = \alpha_3 e_3 + \alpha_4 e_4$. Then we can write

$$e_1 \circ e_1 = \alpha_2 e_4, \quad e_1 \circ e_2 = e_3, \quad e_2 \circ e_1 = \alpha_5 e_3 + \alpha_6 e_4,$$

$$e_2 \circ e_2 = \alpha_8 e_4, \quad e_1 \circ e_3 = e_4, \quad e_2 \circ e_3 = \alpha_9 e_4.$$

Consider the products

$$(e_1 \circ e_2) \circ e_1 = e_1 \circ (e_2 \circ e_1) + e_1 \circ (e_1 \circ e_2) = e_1 \circ (-e_3 + \alpha_6 e_4) + e_1 \circ e_3 = 0 \Rightarrow e_3 \circ e_1 = 0,$$

$$(e_1 \circ e_2) \circ e_2 = 2e_1 \circ (e_2 \circ e_2) = 0 \Rightarrow e_3 \circ e_2 = 0.$$

If we replace basic elements as follows:

$$e'_1 = e_1, \quad e'_2 = e_2 - \alpha_9 e_1, \quad e'_3 = e_3 - \alpha_2 \alpha_9 e_4, \quad e'_4 = e_4,$$

we obtain $\alpha_9 = 0$, i.e. the multiplication in the algebra has the following form:

$$e_1 \circ e_1 = \alpha e_4, \quad e_1 \circ e_2 = e_3,$$

$$e_2 \circ e_1 = -e_3 + \beta e_4, \quad e_2 \circ e_2 = \gamma e_4, \quad e_1 \circ e_3 = e_4$$

(omitted products are equal to zero).

Check the isomorphism inside this family.

Consider the general change of generator basic elements:

$$e'_1 = a_1 e_1 + a_2 e_2 + a_3 e_3,$$

$$e'_2 = b_1 e_1 + b_2 e_2 + b_3 e_3$$

where $a_1 b_2 - a_2 b_1 \neq 0$.

Expressing rest basic elements of the new basis via basic elements of the old basis and comparing the obtained relations we obtain the following restrictions:

$$a_1^2 \alpha + a_1 a_2 \beta + a_2^2 \gamma + a_1 a_3 = \alpha' a_1^2 b_2,$$

$$a_1 b_2 \beta + 2 a_2 b_2 \gamma + a_1 b_3 = \beta' a_1^2 b_2,$$

$$b_2 \gamma = \gamma' a_1^2, \quad b_1 = 0.$$

Consider the following cases

Case 2.1. Let $\gamma = 0$. Then $\gamma' = 0$ and

$$a_1 \alpha + a_2 \beta + a_3 = \alpha' a_1 b_2,$$

$$b_2 + b_3 = \beta' a_1 b_2.$$

Taking $a_3 = -a_1 \alpha - a_2 \beta$ and $b_3 = -b_2$ we obtain $\alpha' = \beta' = 0$, i.e. we have the algebra a_4 .

Case 2.2. Let $\gamma \neq 0$. Then putting $b_2 = \frac{a_1^2}{\gamma}$, $a_3 = -\frac{a_1^2 \alpha + a_1 a_2 \beta + a_2^2 \gamma}{a_1}$, and $b_3 = \frac{a_1 b_2 \beta + 2 a_2 b_2 \gamma}{a_1}$, we get $\gamma' = 1$, $\alpha' = \beta' = 0$, i.e. the algebra a_5 is obtained.

Note that algebras with the conditions $(2, 0, 0)$, $(1, 0, 0)$, $(0, 0, 0)$ are associative. Therefore we can use the classification of four dimensional algebras of Leibniz [1], i.e. choose algebras with the condition $A^3 = 0$. $\square$

References

- [1] S. Albeverio, B.A. Omirov and I.S. Rakhimov, Classification of 4-dimensional nilpotent complex Leibniz algebras. arXiv.math.RA/0611831 (accepted in Extracta Mathematicae).
- [2] Sh.A. Ayupov and B.A. Omirov, On some classes of nilpotent Leibniz algebras. Sib. Math. J. vol. 42(1) (2001), 18-29.
- [3] A.S. Dzhumadil'daev, Identities for multiplications derived by Leibniz and Zinbiel multiplications. Abstracts of short communications of International conference "Operator algebras and quantum theory of probability" (2005), Tashkent, 76-77.

- [4] A.S. Dzhumadil'daev and K.M. Tulenbaev, Nilpotency of Zinbiel algebras. J. Dyn. Control. Syst., vol. 11(2) (2005), 195-213.
- [5] J.-L. Loday, Cup product for Leibniz cohomology and dual Leibniz algebras. Math Scand., 77 (1995), 189-196.
- [6] J.-L. Loday and T. Pirashvili, Universal enveloping algebras of Leibniz algebras and (co)homology. Math. Ann. 296 (1993), 139-158.
- [7] J.-L. Loday, A. Frabetti, F. Chapoton and F. Goichot, Dialgebras and Related Operads. Lecture Notes in Math., vol. 1763, 2001.
- [8] M. Vergne, Cohomologie des algebras de Lie nilpotents. Application a l'etude de la variete des algebres Lie nilpotents. Bull. Soc. Math. France. vol. 98 (1970), 81-116.